

Section Drugs

Drugs given before cardiac arrest

For each drug given, indicate whether given at induction, during maintenance phase or in recovery - do not include treatment c
arrest

1

Grouping (Pick One)

- Anaesthetic agents
- Analgesics and adjuncts
- Neuromuscular Blockade
- Neuromuscular Reversal
- Inotropes/Vasopressors
- Antibiotics
- Antiemetics
- IV Fluids
- Blood products
- Coagulation Products
- Antidotes/reversal agents
- Other adjuncts
- Other exposures and high-risk allergens

2

Drug given (Pick One)

- Propofol
- Thiopental
- Etomidate
- Midazolam
- Ketamine
- Sevoflurane
- Desflurane
- Isoflurane
- Nitrous oxide
- Paracetamol
- Morphine
- Diamorphine
- Fentanyl
- Alfentanil
- Remifentanil
- Codeine/Dihydrocodeine
- Oxycodone
- Tramadol
- Methadone
- Buprenorphine
- NSAIDs
- Gabapentinoid
- IV lidocaine
- Clonidine
- Dexmedetomidine
- Ketamine
- Suxamethonium
- Atracurium/cis-atracurium
- Rocuronium
- Mivacurium
- Vecuronium
- Pancuronium
- Neostigmine + anticholinergic
- Sugammadex
- Ephedrine
- Metaraminol
- Phenylephrine
- Adrenaline
- Noradrenaline
- Dobutamine
- Dopamine
- Dopexamine
- Enoximone
- Glucagon
- Isoprenaline
- Methylene blue
- Milrinone
- Terlipressin
- Vasopressin
- Penicillins

- Cephalosporins
- Metronidazole
- Teicoplanin
- Gentamicin
- Vancomycin
- Ondansetron
- Dexamethasone
- Cyclizine
- Prochlorperazine
- Metoclopramide
- Droperidol
- Crystalloid
- Gelatin or gelatin-containing
- Starch or starch-containing
- Albumin
- Red cells
- Cell salvage blood
- Platelets
- Fresh frozen plasma
- Cryoprecipitate
- Fibrinogen concentrate
- Heparin (any)
- Tranexamic acid
- Aprotinin
- Protamine
- Vitamin K
- Prothrombin complex concentrate
- Recombinant Factor VIIa
- Naloxone
- Flumazenil
- Doxapram
- Magnesium sulphate
- Calcium chloride/gluconate
- Uterotonics
- Patent blue dye
- Methylene blue dye
- Bone cement
- Radio-opaque contrast
- Chlorhexidine
- Other

3

Induction (Pick One)

- Yes
- No

4

Maintenance (Pick One)

- Yes
- No

5

Recovery (Pick One)

- Yes
- No

6

If Yes to blood products:
Emergency O +/- blood
(True / False)

Was patient cross-matched pre-op?
(True / False)

Blood/fluid warmer
(True / False)

Rapid infusion device (e.g. Level 1)
(True / False)

Major haemorrhage protocol activated
(True / False)

None of the above
(True / False)

Section Blocks

Regional/local anaesthesia

1

Please specify the block performed

Grouping (Pick One)

- Upper limb/neck
- Lower limb
- Trunk
- Neuraxial
- Other
- None

2

Block performed (Pick One)

- Interscalene
- Supraclavicular
- Infraclavicular
- Axillary
- Cervical Plexus Deep
- Cervical Plexus Combined
- Cervical Plexus Superficial
- Peripheral nerve(s)
- 3 in 1
- Adductor Canal
- Ankle
- Fascia Iliaca
- Femoral
- Lumbar Plexus
- Popliteal
- Sciatic
- Peripheral nerve(s)
- Erector Spinae
- Intercostal
- Interpleural
- Paravertebral
- Pectoralis
- Quadratus Lumborum
- Rectus Sheath
- Serratus Anterior
- Serratus Plane
- Transversus Abdominus Plane
- Ophthalmic
- Genicular
- Ilioinguinal
- Penile
- Pudendal
- Scalp
- Epidural
- Epidural top-up
- CSE
- Spinal
- Caudal
- Local infiltration
- Biers block
- Other

Please specify

3

Performed by (Pick One)

- Anaesthetist
- Surgeon

4

Landmark (Pick One)

Yes

No

5

Ultrasound (Pick One)

Yes

No

6

Nerve stimulator (Pick One)

Yes

No

7

Catheter placed (Pick One)

Yes

No

8

Drug(s)

9

Concentration(s)

10

Volume(s)

Section 04

Drugs given before cardiac arrest

4.1

Was drug dosing considered to contribute to cardiac arrest? (Pick One)

Yes

No

Please specify

4.2

Drugs given before cardiac arrest

Please Fill in: **Section Drugs**

Multiple copies may be included.

Regional/local anaesthesia

4.3

Was Local anaesthetic given before cardiac arrest (Pick One)

Yes

No

4.4

Regional/local anaesthesia

Please Fill in: **Section Blocks**

Multiple copies may be included.